

# mimi.

Medication  
Abortion Kit  
(MIFEPRISTONE  
+ MISOPROSTOL)

## Drug Facts (continued)

### When using this product

**Most people will have these symptoms**, which are normal and usually a sign the medications are working.

- Cramping that is stronger than a menstrual period (cramping may continue for several days)
- Bleeding that is heavier than a menstrual period (lighter bleeding may continue for up to 2 weeks)
- These symptoms usually last for about a day after taking misoprostol ■:

- Nausea and vomiting
- Diarrhea
- Weakness
- Headache
- Dizziness
- Mild fever or chills

If needed, you can take over-the-counter medications for these symptoms; see the leaflet inside the package for additional information.

**Talk with a health care provider or seek medical help** if you have any of these rare symptoms, which may be a sign of something serious such as ectopic pregnancy, infection, or too much bleeding.

- Severe pain that does not get better with pain medicine
- Bleeding that soaks through 2 full-size pads per hour for 2 consecutive hours
- No bleeding or only light bleeding within 2 days of taking misoprostol ■
- "Feeling sick" with several symptoms, including continued pain, nausea, vomiting, diarrhea, or weakness more than 24 hours after taking misoprostol ■
- Fever of 100.4°F (38°C) or higher that lasts for more than 4 hours

See the leaflet inside the package for additional information.

### This treatment does not always work.

- About 3 out of 100 users of this treatment will need an in-clinic procedure (vacuum aspiration) to complete the abortion.

## Drug Facts (continued)

### Directions

**Step 1:** Swallow 1 mifepristone ● pill.

**Step 2** (24-48 hours after taking mifepristone ●):

**2a:** Place 4 misoprostol ■ pills between your gums and cheeks (2 on the bottom left side of your mouth, and 2 on the bottom right side). Keep the tablets there for up to 30 minutes while they dissolve, then swallow anything that remains with water.

**2b:** 4 hours later, take the additional 4 misoprostol ■ pills if you:

- are more than 8 weeks pregnant, OR
- have had no bleeding or only light bleeding

To take the additional 4 misoprostol ■ pills, see Step 2a.

**Step 3:** Monitor your symptoms.

**Step 4** (4 weeks after taking mifepristone ●): Take a urine pregnancy test. If the test is positive, you should contact a health care provider to discuss treatment options. See the leaflet inside the package for additional information.

### Other information

Store at room temperature 20°C - 25°C (68°F - 77°F).

### Inactive ingredients

Mifepristone ●: colloidal silica anhydrous, corn starch, magnesium stearate, microcrystalline cellulose, and povidone.

Misoprostol ■: hydrogenated castor oil, hypromellose, microcrystalline cellulose, and sodium starch glycolate.

### Questions or comments?

If you have questions, please contact the clinic where you are receiving care.



Read entire leaflet inside the package before beginning use of this product. Scan QR code for more information.  
[www.mimi-pills.com](http://www.mimi-pills.com)

Manufactured for  
Advancing New Standards in Reproductive Health  
Oakland, CA

## Medication Abortion Kit (MIFEPRISTONE + MISOPROSTOL)

Contains  
1 200 mg Mifepristone tablet  
8 200 mcg Misoprostol tablets  
1 urine pregnancy test

## Drug Facts

### Active ingredients (in each tablet)

Mifepristone ● 200 mg (1 pill).....blocks a hormone needed for pregnancy to continue  
Misoprostol ■ 200 mcg (8 pills).....causes pregnancy to be passed from uterus

### Purpose

### Use

Use **both** medicines separately as directed to cause an abortion in early pregnancy (up to 10 weeks since the first day of your last menstrual period)

### Warnings

**Bleeding and infection alert:** Heavy bleeding and serious infections may occur following childbirth, miscarriage, or abortion, including when using this product. These can be fatal. It is important to seek medical help if you have heavy bleeding or signs of infection (see "When using this product").

### Allergy alert:

Do not use if you have ever had an allergic reaction to mifepristone ● or misoprostol ■.

### Do not use if you:

- are more than 10 weeks (70 days) pregnant
- since you got pregnant, you have missed more than 2 periods
- got pregnant while using an IUD (intrauterine device)
- take a medicine to thin your blood (anticoagulant)
- take steroid pills (such as prednisone)
- have ever had, or currently have, a pregnancy outside the uterus (ectopic pregnancy)
- have had surgery on your fallopian tubes, or had your tubes tied
- have been told by a doctor or other health care provider that you have:
  - a bleeding disorder (a condition where your blood does not clot normally)
  - problems with your adrenal glands (chronic adrenal failure)
  - a disease called porphyria

### Ask a doctor or other health care provider before use if you

- are unsure if you are less than 10 weeks pregnant
- have had a blood transfusion for severe anemia in the past year
- have had unexplained bleeding during this pregnancy
- want more information about your options for continuing or ending this pregnancy